

Ground-Based Techniques in Mangrove Monitoring and Management

1
2

**Sharif A. Mukul, Mohammed Abu Sayed Arfin-Khan, Saon Banerjee,
Dhananjay Barman, Gouranga Kar, Prashant K. Srivastava,
Pallab Mozumder, and Sanjeev Kumar Srivastava**

3
4
5

Abstract Mangroves provide a wide range of ecosystem services, including coastal protection, biodiversity conservation, and fisheries support. While offering cultural and recreational benefits to local communities, they also play a crucial role in climate regulation and nutrient cycling. Despite their immense importance, mangrove ecosystems around the world are disappearing at an alarming rate. To safeguard this critical coastal ecosystem, effective management and continuous monitoring are

6
7
8
9
10
11

S. A. Mukul (✉)

Department of Environment and Development Studies, United International University,
Dhaka, Bangladesh

School of Science, Technology and Engineering (SSTE), University of the Sunshine Coast,
Sippy Downs, QLD, Australia

Department of Earth and Environment, Florida International University, Miami, FL, USA
e-mail: mukul@eds.uui.ac.bd

M. A. S. Arfin-Khan

Department of Forestry and Environmental Science, Shahjalal University of Science and
Technology, Sylhet, Bangladesh

S. Banerjee

Department of Agricultural Meteorology and Physics, Bidhan Chandra Krishi
Viswavidyalaya, Mohanpur, West Bengal, India

D. Barman · G. Kar

ICAR—Central Research Institute for Jute and Allied Fibers, Kolkata, West Bengal, India

P. K. Srivastava

School of Science, Technology and Engineering (SSTE), University of the Sunshine Coast,
Sippy Downs, QLD, Australia

Institute of Environment and Sustainable Development, Banaras Hindu University,
Varanasi, Uttar Pradesh, India

P. Mozumder

Department of Earth and Environment, Florida International University, Miami, FL, USA

S. K. Srivastava

School of Science, Technology and Engineering (SSTE), University of the Sunshine Coast,
Sippy Downs, QLD, Australia

12 essential. Ground-based monitoring techniques play a crucial role by providing
13 high-resolution, site-specific data that complement remote sensing technologies,
14 offering detailed insights into mangrove health, dynamics, and resilience. This
15 chapter presents an overview of ground-based approaches to mangrove manage-
16 ment and monitoring, emphasizing their significance; the key biological, physical,
17 chemical, and social parameters involved; important methodological consider-
18 ations; and the challenges associated with their application. While ground-based
19 monitoring faces challenges such as limited accessibility, high resource demands,
20 and environmental risks, it remains indispensable for the long-term conservation
21 and management of mangroves. By combining these techniques with community
22 engagement and technological innovation, stakeholders can develop robust strate-
23 gies to strengthen mangrove resilience and ensure the sustainable management and
24 protection of this imperiled ecosystem.

25 **Keywords** Biogeochemical process · Climate change · Ecosystem services ·
26 Livelihood dependence · Mangrove forest

27 1 Introduction

28 Mangroves are coastal ecosystems that occur in the intertidal zones of tropical and
29 subtropical regions worldwide (Fig. 1; Spalding et al., 2010). Globally, they cover
30 about 15 million hectares across 118 countries, accounting for nearly 15% of the
31 world’s coastlines (IUCN, 2024; Giri et al., 2011; Table 1). Mangrove ecosystems
32 are characterized by the presence of “true mangroves,” i.e., plant species that are
33 specially adapted to saline environments, featuring unique traits such as



Fig. 1 Map showing the global distribution of mangroves (in black). Source: UNEP-WCMC (<https://data.unep-wcmc.org>; accessed on 15 June 2024)

Table 1 Mangrove distribution by region

Region	Share of global mangroves (%)	Major countries
Asia	42	Indonesia, Malaysia, Thailand
Africa	21	Mozambique, Kenya, Tanzania
North and Central America	15	Mexico, Belize, USA (Florida)
Australia and Oceania	12	Australia, Papua New Guinea
South America	10	Brazil, Colombia, Venezuela

pneumatophores (aerial roots), prop roots, and salt-excreting leaves (Biswas & Biswas, 2019; Lugo & Snedaker, 1974, Table 2).

Mangrove ecosystems provide a wide range of critical ecosystem services (Ho & Mukul, 2021). They act as natural barriers, protecting coastlines from cyclones, tsunamis, and storm surges (Mohammed et al., 2024; Costanza et al., 2008). They also play a significant role in carbon storage and sequestration (Hamilton & Friess, 2018; Donato et al., 2011), contribute to water quality regulation (Romañach et al., 2018), and serve as critical breeding and nursery habitats for numerous fish and wildlife species (zu Ermgassen et al., 2025). Beyond their ecological functions, mangroves also support local livelihoods by providing resources such as food, fuel, timber, construction materials, and opportunities for ecotourism (Friess et al., 2019).

Despite their immense ecological and socio-economic importance, mangroves are disappearing at a rate faster than that of tropical rainforests (Thomas et al., 2017). According to the first global assessment of mangroves under the IUCN Red List of Ecosystems, more than half of the world's mangrove ecosystems are currently at risk of collapse (Fig. 2; IUCN, 2024). The primary drivers of mangrove loss include coastal infrastructure development, aquaculture, conversion of land for agriculture, climate change, and sea-level rise (Leal & Spalding, 2024; Mukul et al., 2019, 2020a). Between 2000 and 2016, over half of global mangrove loss was attributed to land-use change, mainly the conversion of mangrove areas into aquaculture and agricultural lands, with the majority of these losses concentrated in six Southeast Asian countries (FAO, 2023; Goldberg et al., 2020).

Given their critical role in carbon sequestration, coastal protection, biodiversity conservation, and community livelihoods, the sustainable management and effective monitoring of mangroves are imperative to safeguard these vital ecosystems from ongoing and future threats (Friess et al., 2024; Schmitt & Duke, 2015). Ground-based techniques are essential for providing detailed, localized, and highly accurate data for the assessment and management of mangroves and for complementing remote sensing and aerial data (Misiukas et al., 2021).

This chapter provides an overview of the rationale for mangrove management and monitoring, emphasizing the importance of ground-based approaches. It discusses the key biological, physical, and chemical processes that are essential for effective mangrove monitoring and management, including human dimensions, main methodological considerations, and the challenges associated with ground-based monitoring and management of mangroves.

Table 2 List of true mangrove species

Family	Species
Acanthaceae	<i>Avicennia alba</i>
	<i>Avicennia germinans</i> (black mangrove)
	<i>Avicennia marina</i>
	<i>Avicennia officinalis</i>
Combretaceae	<i>Laguncularia racemosa</i> (white mangrove)
	<i>Lumnitzera littorea</i>
	<i>Lumnitzera racemosa</i>
Lythraceae	<i>Sonneratia alba</i>
	<i>Sonneratia apetala</i>
	<i>Sonneratia caseolaris</i>
	<i>Sonneratia griffithii</i>
	<i>Sonneratia ovata</i>
Rhizophoraceae	<i>Bruguiera cylindrica</i>
	<i>Bruguiera exaristata</i>
	<i>Bruguiera gymnorhiza</i>
	<i>Bruguiera hainesii</i>
	<i>Bruguiera parviflora</i>
	<i>Bruguiera sexangula</i>
	<i>Ceriops australis</i>
	<i>Ceriops decandra</i>
	<i>Ceriops tagal</i>
	<i>Kandelia candel</i>
	<i>Kandelia obovata</i>
	<i>Rhizophora apiculata</i>
	<i>Rhizophora harrisonii</i>
	<i>Rhizophora mangle</i> (red mangrove)
	<i>Rhizophora mucronata</i>
	<i>Rhizophora racemosa</i>
<i>Rhizophora stylosa</i>	
Arecaceae (Palms)	<i>Nypa fruticans</i> (special mangrove palm)
Malvaceae	<i>Heritiera fomes</i>
	<i>Heritiera littoralis</i>
	<i>Hibiscus tiliaceus</i>
Rubiaceae	<i>Scyphiphora hydrophylacea</i>
Euphorbiaceae	<i>Excoecaria agallocha</i>
	<i>Excoecaria indica</i>
Plumbaginaceae	<i>Aegialitis annulata</i>
	<i>Aegialitis rotundifolia</i>




Fig. 2 Global red list of mangrove ecosystems. Source: Modified after IUCN (2024)



Fig. 3 Vertical (left) and horizontal (right) representation of a mangrove ecosystem. Photo credits: Sharif A. Mukul

2 The Rationale for Mangrove Management and Monitoring

69

The objective of mangrove management typically falls into several key categories: sustainable timber production, conservation of forests and wildlife habitats, coastal and shoreline protection, ecological restoration, and support of essential ecological functions such as soil stabilization, site protection, and carbon sequestration (Akram et al., 2023). The rationale for managing mangroves may also vary depending on the position or type of mangrove forest (Schmitt & Duke, 2015). For instance, as shown in Fig. 3, fringe mangrove forests play a vital role in protecting shorelines, and riverine forests are especially important for supporting plant and animal productivity,

70
71
72
73
74
75
76
77

78 while basin or interior mangroves serve as key nutrient sinks and sources of wood
79 products (Ewel et al., 1998).

80 Objectives of mangrove management change over time and may include specific
81 goals such as coastal protection through coastal green belt or offsetting carbon
82 emissions (Schmitt & Duke, 2015). Historically, the primary objective was on affor-
83 estation for silviculture, emphasizing coastal stabilization and timber production to
84 support industries like pulp and paper manufacturing (Iftekhar & Islam, 2004). Over
85 time, greater attention shifted toward the ecological functions of mangroves, includ-
86 ing their role as wildlife habitats, coastal protection, contributors to pelagic food
87 webs, and providers of livelihoods for coastal communities (Friess et al., 2019;
88 Stubbs & Saenger, 2002). More recently, mangroves have been managed as cost-
89 effective, nature-based solutions for enhancing climate resilience, acknowledged
90 for their capacity to sequester atmospheric carbon more rapidly than other tropical
91 ecosystems and their capacity to endure rising sea levels (Mukul et al., 2020b;
92 Krauss et al., 2014).

93 Mangroves grow in different hydrogeomorphic settings, such as river-, tide-,
94 wave-dominated, or interior mangrove forests, and rainfall, temperature, and fresh-
95 water supply also affect mangrove growth (Biswas & Biswas, 2019). Figure 4 pres-
96 ents a generalized decision-making flowchart that can guide mangrove management
97 processes. The primary objective of any mangrove management strategy should be
98 to protect and conserve existing mangrove ecosystems while minimizing the distur-
99 bances driving their loss. This approach is often more efficient and more cost-
100 effective than efforts focused solely on planting new mangroves (Leal & Spalding,
101 2024). Alongside legal protection, local communities play a vital role in safeguard-
102 ing mangrove forests from degradation and loss (Datta et al., 2012). Raising aware-
103 ness and actively involving local stakeholders in mangrove management can further
104 strengthen conservation outcomes (Walters et al., 2008).

105 In areas where mangroves have been present in the past, it is important to distin-
106 guish between sites that have been degraded since the loss of mangrove cover and
107 those that remain relatively intact. Based on this distinction and having determined
108 that planting is necessary, three different types of mangrove planting can be
109 employed: reforestation, rehabilitation, and restoration (Schmitt & Duke, 2015).
110 Additionally, afforestation is used in areas where mangroves did not previously
111 grow (Siddiqi, 2001).

112 Because species distribution is closely tied to specific site conditions, careful
113 attention must be given first to selecting a suitable planting site, followed by choos-
114 ing the appropriate species and the most effective planting technique for that loca-
115 tion (Lewis & Brown, 2014). Planting can be carried out using seeds, propagules, or
116 seedlings, and site selection should be guided by an analysis of historical changes
117 and natural processes (Schmitt & Duke, 2015). Historical data helps clarify coastal
118 dynamics, such as patterns of accretion and erosion, and supports the selection of
119 species that naturally thrived in the area prior to human disturbance (Lewis &
120 Brown, 2014). This process should be complemented by observing natural regen-
121 eration, which signals that a site is suitable for mangrove growth and provides valu-
122 able insights into appropriate species choices and planting methods (Siddiqi, 2001).




Fig. 4 Decision-making flow chart for mangrove management. Source: Modified after Schmitt and Duke (2015)

Mangrove forests have a strong capacity for natural regeneration following disturbances (Siddiqi, 2001). In areas where mangroves have been destroyed or degraded, a thorough site assessment is required to determine whether interventions such as soil or hydrological restoration are necessary before planting (Lewis & Brown, 2014). Whenever possible, rehabilitation approaches that promote natural regeneration should be prioritized (Schmitt & Duke, 2015). However, in cases where habitat loss or degradation is so severe that natural processes are insufficient for recovery, tailored, site-specific, and cost-effective rehabilitation or restoration methods must be implemented (Lewis & Brown, 2014).

Large-scale afforestation is typically undertaken to establish mangrove forests on treeless mudflats or newly accreted lands (Siddiqi, 2001). In such areas,

134 afforestation requires specialized and carefully designed planting techniques. After
135 planting, seedlings must be protected from human disturbances, such as destructive
136 fishing practices and grazing by cattle or sheep (Schmitt & Duke, 2015). In some
137 locations, additional protection from wave action may also be necessary. Once man-
138 groves are established, they must be effectively managed and safeguarded from
139 human activities such as felling or conversion to other land uses. Traditionally, these
140 management and protection efforts have been overseen by government forestry
141 departments.

142 Mangrove monitoring refers to the systematic collection of data and the process-
143 ing of this data into information about the condition, health, and status of mangrove
144 forests (Schmitt & Duke, 2015). It also helps to understand why changes are occur-
145 ring. Ground-based monitoring of mangroves involves direct, on-site assessments
146 of this ecosystem using appropriate approaches. While recent technological
147 advancements, such as the availability of satellite imagery, artificial intelligence,
148 remote sensing, and machine learning, have provided new opportunities for moni-
149 toring mangroves, ground-based data remains essential (Wang et al., 2025).

150 **3 Considerations for Ground-Based Management** 151 **and Monitoring of Mangroves**

152 Mangrove ecosystems are shaped by a complex interplay of biological, physical,
153 and chemical processes that sustain their structure, function, and resilience (Lugo &
154 Snedaker, 1974). The ecological integrity of these ecosystems is maintained through
155 the synergistic interaction of these processes, which operate across various spatial
156 and temporal scales (Fig. 5). In addition to these natural dynamics, human dimen-
157 sions—including anthropogenic pressures such as disturbances and local depen-
158 dence on mangrove resources—play a significant role in influencing mangrove
159 health and stability (Biswas et al., 2009). Disruption of any of these interconnected
160 components, whether ecological or human-driven, can compromise the functional-
161 ity of mangroves and the vital ecosystem services they provide (Schmitt & Duke,
162 2015). Therefore, a comprehensive understanding of these processes is essential for
163 the effective monitoring and management of mangrove ecosystems, as outlined in
164 the sections that follow.

165 **3.1 Biological Processes**

166 Biological processes in mangrove ecosystems are essential for sustaining productiv-
167 ity, maintaining biodiversity, and ensuring resilience to environmental change (Lugo
168 & Snedaker, 1974). Primary production, driven by mangrove trees, algae, and epi-
169 phytic organisms, supports both local consumers and the export of energy and




Fig. 5 The interplay of physical, chemical, and biological processes in the mangrove ecosystem

nutrients to adjacent ecosystems (Alongi, 2002). Litterfall and detrital pathways provide organic material that fuels microbial decomposition and detritivore food webs (Twilley et al., 1992). Faunal interactions and bioturbation, carried out by organisms such as crabs, mollusks, and polychaetes, play crucial roles in sediment aeration, nutrient cycling, and organic matter turnover (Kristensen, 2008). Their burrowing activities enhance oxygen penetration and stimulate microbial processes in otherwise anoxic sediments. Symbiotic relationships, such as nitrogen-fixing bacteria associated with mangrove roots, further contribute to nitrogen inputs in these typically nutrient-poor coastal environments (Kristensen et al., 2008). Finally, mangrove regeneration is shaped by recruitment, seedling establishment, and species interactions, which together drive successional dynamics and determine zonation patterns within mangrove forests (Siddiqi, 2001).

3.2 Physical Processes

Physical processes in mangroves shape their zonation patterns, habitat formation, and interactions with neighboring ecosystems such as seagrass beds and coral reefs. These processes are largely driven by hydrodynamics, geomorphology, and climatic forces. For example, tidal action regulates inundation patterns, sediment deposition, and nutrient fluxes across the intertidal zone (Alongi, 2009). Tidal exchange also plays a crucial role in oxygenating mangrove soils and maintaining salinity

189 gradients within the water column. Meanwhile, sediment dynamics in mangroves
190 enhance sediment trapping and vertical accretion, contributing to land-building pro-
191 cesses that help buffer against sea-level rise (Krauss et al., 2014). The dense root
192 networks of mangroves attenuate wave energy, stabilize shorelines, reduce coastal
193 erosion, and promote further sediment deposition (Mazda et al., 1997). Additionally,
194 storms and cyclonic disturbances can reshape mangrove stands by altering sediment
195 profiles and clearing vegetation, triggering successional processes that influence
196 ecosystem recovery (Duke, 2001).

197 **3.3 Chemical Processes**

198 Chemical processes in mangrove ecosystems are largely shaped by anaerobic soil
199 conditions and tidal influences, playing a central role in the mangroves' contribu-
200 tion to global biogeochemical cycles (Kristensen et al., 2008). Mangroves help
201 regulate salinity through specialized physiological adaptations, such as salt exclu-
202 sion at the roots, salt secretion via specialized glands, or salt storage within vacuoles
203 (Lugo & Snedaker, 1974). The sediments in mangrove systems are typically anoxic,
204 fostering reduction reactions like sulfate reduction, methanogenesis, and denitrifi-
205 cation, all of which influence nutrient dynamics and availability (Alongi, 1994). In
206 addition, nitrogen cycling—including ammonification, nitrification, and denitrifica-
207 tion—occurs within both the sediments and rhizosphere, further regulating nitrogen
208 availability (Kristensen et al., 2008). Phosphorus cycling is primarily governed by
209 adsorption-desorption processes involving iron and aluminum oxides in the sedi-
210 ments. Another key chemical process is the decomposition of organic matter, where
211 mangrove leaf litter and woody debris break down through microbial activity,
212 releasing dissolved organic carbon and nutrients that sustain detrital food webs
213 (Alongi, 2002). Importantly, mangrove soils act as major carbon sinks, with the
214 long-term burial of organic carbon under anaerobic conditions significantly contrib-
215 uting to “blue carbon” storage (Mukul et al., 2020b).

216 **3.4 Human Dimensions**

217 The human dimensions of mangroves encompass the complex social, cultural, and
218 economic relationships between coastal communities and these vital ecosystems
219 (Biswas et al., 2009). Mangroves provide a diverse array of ecosystem services,
220 including fisheries, timber, fuelwood, non-timber forest products (NTFPs), coastal
221 protection, and ecotourism opportunities, upon which millions of people worldwide
222 depend for their livelihoods and well-being (Sarkar et al., 2024; Fig. 6). Therefore,
223 the effective management and monitoring of mangroves must account for socio-
224 economic factors and actively engage local communities in decision-making to
225 ensure both ecological sustainability and community benefits (Datta et al., 2012).



Fig. 6 Mangroves provide key non-timber products like *Nypa* palm (left) and support ecotourism (right), as seen in the Sundarbans forest, where both generate important revenue and showcase the ecosystem’s rich biodiversity. Photo credits: Sharif A. Mukul

4 Ground-Based Approaches for Monitoring and Managing Mangroves

226
227

Ground-based approaches play a crucial role in effective mangrove monitoring and management by providing detailed, site-specific information. These approaches include field-based surveys to assess forest structure, species composition, biomass, and regeneration status, as well as monitoring of soil and water parameters such as salinity, nutrient levels, and sedimentation (Bosire et al., 2008; Lugo & Snedaker, 1974). Participatory monitoring, involving local communities and stakeholders, enhances the accuracy and relevance of data collection by integrating traditional ecological knowledge and local observations (Walters et al., 2008). Ground-based methods also support adaptive management by enabling regular assessments of human pressures, such as harvesting intensity, land-use changes, and pollution, which are critical for designing locally appropriate conservation and restoration strategies (Datta et al., 2012). When combined with socio-economic data, these field-based approaches provide a holistic framework for managing mangroves in ways that balance ecological health with community needs (Biswas et al., 2009).

Ground-based techniques are also essential for validating and calibrating remote sensing data. Field measurements—such as vegetation density, species identification, and canopy structure—provide critical ground-truth information that improves the accuracy of satellite or aerial imagery analysis (Maurya et al., 2021). By integrating ground-based and remote sensing approaches, the overall effectiveness and precision of mangrove monitoring have significantly improved (Karsch et al., 2023). Recent technological advancements have further improved ground-based monitoring, with tools like handheld GPS units, drones, and mobile apps enabling more efficient data collection and real-time reporting. For instance, drones can capture high-resolution images of mangrove areas, while mobile apps allow field teams to record and upload data directly to centralized databases, streamlining both analysis and management efforts.

228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253

254 **4.1 Field-Based Surveys and Sampling**

255 Field surveys and sampling are the cornerstone of ground-based mangrove monitoring
 256 and management, providing essential, on-the-ground data. These activities
 257 involve direct observations and measurements of vegetation, fauna, soil, water
 258 parameters, and biogeochemical processes (Fig. 7). Key components include:

- 259 • **Biodiversity surveys:** Recording the presence and abundance of mangrove flora
 260 and fauna—including fish, crabs, birds, predators, and prey—to assess species
 261 composition, diversity, and ecological interactions (Mukul et al., 2019).
- 262 • **Vegetation assessment:** Measuring tree height, diameter at breast height (DBH),
 263 species composition, and canopy cover from representative plots of different size
 264 and shape to evaluate mangrove forest health, biomass, and carbon storage
 265 potential (Mukul et al., 2020b).
- 266 • **Seedling recruitment:** Monitoring the density and distribution of mangrove
 267 seedlings to gauge regeneration potential and forest recovery capacity
 268 (Siddiqi, 2001).
- 269 • **Phenology monitoring:** Tracking seasonal changes in flowering, fruiting, and
 270 leaf development to better understand mangrove life cycles and reproductive pat-
 271 terns in the face of climate or salinity change (Siddiqi, 2001).
- 272 • **Soil sampling:** Analyzing soil properties such as salinity, pH, organic matter
 273 content, and nutrient levels, which are critical for understanding growth condi-
 274 tions and detecting stressors like pollution or sediment imbalance (Twilley
 275 et al., 1992).
- 276 • **Sedimentation and erosion monitoring:** Using techniques such as sediment
 277 traps, erosion pins, and marker horizons to measure sediment deposition and



Fig. 7 Snapshots from field surveys conducted in the Sundarbans mangrove forest of Bangladesh. Photo credits: Sharif A. Mukul

erosion rates, helping assess the effects of coastal development, sea-level rise, and storm events on mangrove stability (Bhargava & Friess, 2022).

- **Water quality monitoring:** Measuring parameters such as salinity, temperature, dissolved oxygen, turbidity, and nutrient concentrations to understand water conditions influencing mangrove ecosystems (Kristensen et al., 2008). Regular monitoring of water also helps detect pollution sources, including agricultural runoff and industrial discharges, and their impacts on mangrove health.

4.2 Permanent Sample Plots

Establishing permanent plots is a common ground-based technique for long-term mangrove monitoring. These fixed plots are repeatedly surveyed over time to document changes in vegetation structure, species composition, and overall ecosystem dynamics. By tracking key indicators such as growth rates, regeneration, and mortality, permanent plots offer valuable insights into how mangrove forests respond to environmental changes and help predict their future under the influence of drivers such as climate variability, natural disturbances, and species interactions (Misiukas et al., 2021).

4.3 Socio-economic Surveys

Socio-economic surveys explore how mangroves support local livelihoods through traditional and commercial uses, such as harvesting NTFPs, fisheries, and honey production (Sarkar et al., 2024). These surveys help capture the diverse human benefits derived from mangroves and inform management strategies that are tailored to local contexts and community needs (Datta et al., 2012). Mangrove economic value, however, varies depending on valuation methods, the types of ecosystem services assessed, and local socio-economic contexts (Himes-Cornell et al., 2018). Some assessments even use composite indicators like “saved wealth” (avoided damage to property and income) and “saved health” (reduced risk of disease, injury, or death) to estimate the broader adaptation and resilience benefits provided by mangroves (Vo et al., 2012). For instance, coastal mangroves play a critical role in attenuating wave energy, reducing storm surge impacts, and preventing shoreline erosion—services that can be quantified using such approaches (Akber et al., 2018). In addition, mangroves generate substantial revenue through ecotourism, offering opportunities for local employment and environmental education. Socio-economic surveys can offer valuable insights for understanding not only the direct and indirect benefits of ecotourism but also visitors’ willingness to pay for specific ecosystem services, helping guide sustainable tourism development and conservation efforts (Saha & Mukul, 2022).

314 **4.4 Community-Based Monitoring (CBM)**

315 Engaging local communities in mangrove monitoring is a powerful ground-based
316 approach (Walters et al., 2008). Local knowledge and participation enhance data
317 collection efforts and foster a sense of ownership and responsibility for mangrove
318 conservation (Datta et al., 2012). CBM often includes activities like mangrove
319 planting, clean-up drives, and awareness campaigns.

320 **5 Challenges in Ground-Based Mangrove Management** 321 **and Monitoring**

322 Despite its many benefits, ground-based monitoring and management of mangroves
323 face several significant challenges. One of the primary difficulties is accessibility, as
324 mangrove ecosystems are often located in remote, isolated, or difficult-to-reach
325 coastal zones, making fieldwork logistically complex and time-consuming (Halder
326 et al., 2021). These activities are also highly resource-intensive, requiring substan-
327 tial investments of time, labor, specialized skills, and financial resources to conduct
328 surveys, sampling, and data collection effectively. Additionally, field teams working
329 in mangrove environments are often exposed to environmental hazards, including
330 extreme weather, unstable terrain, strong tidal currents, and encounters with poten-
331 tially dangerous wildlife. Security concerns also arise, particularly in remote or
332 less-monitored areas where the presence of armed groups, pirates, or ferocious ani-
333 mals may pose serious threats to field personnel, sometimes necessitating the use of
334 armed guards or specialized protection measures to ensure safe operations.

335 **6 Conclusion**

336 Ground-based techniques are indispensable for effective monitoring and manage-
337 ment of mangrove ecosystems. Despite the rapid advancements in remote sensing
338 technologies, field-based observations provide irreplaceable, fine-scale data on spe-
339 cies composition, forest structure, regeneration status, soil and hydrological condi-
340 tions, and anthropogenic pressures. These techniques—ranging from simple plot
341 surveys and transect measurements to more advanced methods such as soil sam-
342 pling and sediment monitoring—offer critical ground-truthing for interpreting
343 remotely sensed data and are essential for understanding ecological processes that
344 cannot yet be fully captured from above. Furthermore, ground-based approaches
345 play a crucial role in participatory monitoring of mangroves, engaging local com-
346 munities and stakeholders, and integrating traditional ecological knowledge into
347 management practices. This not only improves the accuracy and relevance of moni-
348 toring outcomes but also fosters local stewardship and enhances the sustainability of

conservation efforts in mangroves. Integrating ground-based methods with modern geospatial tools in a complementary, hybrid monitoring framework also holds promise for delivering more comprehensive, multi-scale assessments of mangrove health and change. As mangroves continue to face mounting pressures from climate change and sea-level rise, robust ground-based monitoring will remain a cornerstone of efforts to protect this invaluable coastal ecosystem.

References

- Akber, M. A., Patwary, M. M., Islam, M. A., & Rahman, M. R. (2018). Storm protection service of the Sundarbans mangrove forest, Bangladesh. *Natural Hazards*, *94*, 405–418.
- Akram, H., Hussain, S., Mazumdar, P., Chua, K. O., Butt, T. E., & Harikrishna, J. A. (2023). Mangrove health: A review of functions, threats, and challenges associated with mangrove management practices. *Forests*, *14*, 1698.
- Alongi, D. M. (1994). The role of bacteria in nutrient recycling in tropical mangrove and other coastal benthic ecosystems. *Hydrobiologia*, *285*, 19–32.
- Alongi, D. M. (2002). Present state and future of the world's mangrove forests. *Environmental Conservation*, *29*, 331–349.
- Alongi, D. M. (2009). The energetics of mangrove forests. Springer.
- Bhargava, R., & Friess, D. A. (2022). Previous shoreline dynamics determine future susceptibility to cyclone impact in the Sundarban Mangrove Forest. *Frontiers in Marine Science*, *9*, 814577.
- Biswas, P. L., & Biswas, S. R. (2019). Mangrove forests: Ecology, management, and threats. In W. Leal Filho, A. M. Azul, L. Brandli, A. Lange Salvia, & T. Wall (Eds.), *Life on land. Encyclopedia of the UN Sustainable Development Goals*. Springer.
- Biswas, S. R., Mallik, A. U., Choudhury, J. K., & Nishat, I. (2009). A unified framework for the restoration of Southeast Asian mangroves—Bridging ecology, society and economics. *Wetlands Ecology and Management*, *17*, 365–383.
- Bosire, J. O., Dahdouh-Guebas, F., Walton, M., Crona, B. I., Lewis, R. R., Field, C., Kairo, J. G., & Koedam, N. (2008). Functionality of restored mangroves: A review. *Aquatic Botany*, *89*, 251–259.
- Costanza, R., Pérez-Maqueo, O., Martinez, M. L., Sutton, P., Anderson, S. J., & Mulder, K. (2008). The value of coastal wetlands for hurricane protection. *Ambio*, *37*, 241–248.
- Datta, D., Chattopadhyay, R. N., & Guha, P. (2012). Community based mangrove management: A review on status and sustainability. *Journal of Environmental Management*, *107*, 84–95.
- Donato, D. C., Kauffman, J. B., Murdiyarsa, D., Kurnianto, S., Stidham, M., & Kanninen, M. (2011). Mangroves among the most carbon-rich forests in the tropics. *Nature Geoscience*, *4*, 293–297.
- Duke, N. C. (2001). Gap creation and regenerative processes driving diversity and structure of mangrove ecosystems. *Wetlands Ecology and Management*, *9*, 257–269.
- Ewel, K. C., Twilley, R. R., & Ong, J. E. (1998). Different kinds of mangrove forests provide different goods and services. *Global Ecology and Biogeography Letters*, *7*, 83–94.
- FAO. (2023). *The world's mangroves 2000–2020*. Food and Agriculture Organization of the United Nations (FAO).
- Friess, D. A., Rogers, K., Lovelock, C. E., Krauss, K. W., Hamilton, S. E., Lee, S. Y., Lucas, R., Primavera, J., Rajkaran, A., & Shi, S. (2019). The state of the world's mangrove forests: Past, present, and future. *Annual Review of Environment and Resources*, *44*, 89–115.
- Friess, D. A., Adams, J., Andradi-Brown, D. A., Bhargava, R., Carrasco, G., Dahdouh-Guebas, F., Heck, N., et al. (2024). Mangrove forests: Their status, threats, conservation and restoration. In *Treatise on estuarine and coastal science* (2nd ed.). Elsevier.

- 396 Giri, C., Ochieng, E., Tieszen, L. L., Zhu, Z., Singh, A., Loveland, T., Masek, J., & Duke, N. (2011).
 397 Status and distribution of mangrove forests of the world using earth observation satellite data.
 398 *Global Ecology and Biogeography*, 20, 154–159.
- 399 Goldberg, L., Lagomasino, D., Thomas, N., & Fatoyinbo, T. (2020). Global declines in human-
 400 driven mangrove loss. *Global Change Biology*, 26, 5844–5855.
- 401 Halder, N. K., Merchant, A., Misbahuzzaman, K., Wagner, S., & Mukul, S. A. (2021). Why some
 402 trees are more vulnerable during catastrophic cyclone events in the Sundarbans mangrove for-
 403 est of Bangladesh? *Forest Ecology and Management*, 490, 119117.
- 404 Hamilton, S. E., & Friess, D. A. (2018). Global carbon stocks and potential emissions due to man-
 405 grove deforestation from 2000 to 2012. *Nature Climate Change*, 8, 240–244.
- 406 Himes-Cornell, A., Grose, S. O., & Pendleton, L. (2018). Mangrove ecosystem service val-
 407 ues and methodological approaches to valuation: Where do we stand? *Frontiers in Marine*
 408 *Science*, 5, 376.
- 409 Ho, Y. S., & Mukul, S. A. (2021). Publication performance and trends in mangrove forest: A bib-
 410 liometric analysis. *Sustainability*, 13, 12409.
- 411 Iftekhar, M. S., & Islam, M. R. (2004). Managing mangroves in Bangladesh: A strategy analysis.
 412 *Journal of Coastal Conservation*, 10, 139–146.
- 413 IUCN. (2024). *Red list of mangrove ecosystems*. The World Conservation Union (IUCN). Retrieved
 414 March 25, 2025, from [https://iucn.org/resources/conservation-tool/iucn-red-list-ecosystems/
 415 red-list-mangrove-ecosystems](https://iucn.org/resources/conservation-tool/iucn-red-list-ecosystems/red-list-mangrove-ecosystems)
- 416 Karsch, G., Mukul, S. A., & Srivastava, S. K. (2023). Annual mangrove vegetation cover changes
 417 (2014–2020) in Indian Sundarbans National Park using Landsat-8 and Google Earth Engine.
 418 *Sustainability*, 15, 5592.
- 419 Krauss, K. W., McKee, K. L., Lovelock, C. E., Cahoon, D. R., Saintilan, N., Reef, R., & Chen,
 420 L. (2014). How mangrove forests adjust to rising sea level. *New Phytologist*, 202, 19–34.
- 421 Kristensen, E. (2008). Mangrove crabs as ecosystem engineers; with emphasis on sediment pro-
 422 cesses. *Journal of Sea Research*, 59, 30–43.
- 423 Kristensen, E., Bouillon, S., Dittmar, T., & Marchand, C. (2008). Organic carbon dynamics in
 424 mangrove ecosystems: A review. *Aquatic Botany*, 89, 201–219.
- 425 Leal, M., & Spalding, M. D. (Eds.). (2024). *The state of the world's mangroves 2024*. Global
 426 Mangrove Alliance.
- 427 Lewis, R. R., & Brown, B. (2014). *Ecological mangrove rehabilitation—A field manual for practi-
 428 tioners*. Mangrove Action Project, Canadian International Development Agency and OXFAM.
- 429 Lugo, A. E., & Snedaker, S. C. (1974). The ecology of mangroves. *Annual Review of Ecology and*
 430 *Systematics*, 5, 39–64.
- 431 Maurya, K., Mahajan, S., & Chaube, N. (2021). Remote sensing techniques: Mapping and moni-
 432 toring of mangrove ecosystem—A review. *Complex & Intelligent Systems*, 7, 2797–2818.
- 433 Mazda, Y., Magi, M., Kogo, M., & Nguyen Hong, P. (1997). Mangroves as a coastal protection
 434 from waves in the Tong King delta, Vietnam. *Mangroves and Salt Marshes*, 1, 127–135.
- 435 Misiukas, J. M., Carter, S., & Herold, M. (2021). Tropical forest monitoring: Challenges and
 436 recent progress in research. *Remote Sensing*, 13, 2252.
- 437 Mohammed, S. F., Khan, A., Ahammed, S., Saimun, M. S. R., Bhuiyan, M. S., Srivastava, S. K.,
 438 Mukul, S. A., & Arfin-Khan, M. A. S. (2024). Assessing vulnerability to cyclone hazards in
 439 the world's largest mangrove forest, The Sundarbans: A geospatial analysis. *Forests*, 15, 1722.
- 440 Mukul, S. A., Alamgir, M., Soheli, M. S. I., Pert, P. L., Turton, S. M., Herbohn, J., Khan, M. S. I.,
 441 Ali Reza, A. H. M., Munim, S. A., & Laurance, W. F. (2019). Combined effects of climate
 442 change and sea level rise project dramatic habitat loss of critically endangered Bengal tiger in
 443 the Bangladesh Sundarbans. *Science of the Total Environment*, 663, 830–840.
- 444 Mukul, S. A., Huq, S., Herbohn, J., Seddon, N., & Laurance, W. F. (2020a). Saving the Sundarbans
 445 from development. *Science*, 368, 1198.
- 446 Mukul, S. A., Halim, M. A., & Herbohn, J. (2020b). Forest carbon stock and fluxes: Distribution,
 447 biogeochemical cycles, and measurement techniques. In W. Leal Filho, A. Azul, L. Brandli,

- A. Lange Salvia, & T. Wall (Eds.), *Life on land. Encyclopedia of the UN Sustainable Development Goals*. Springer. 448
449
- Romañach, S. S., DeAngelis, D. L., Koh, H. L., Li, Y., Teh, S. Y., Barizan, R. S. R., & Zhai, L. (2018). Conservation and restoration of mangroves: Global status, perspectives, and prognosis. *Ocean & Coastal Management*, 154, 72–82. 450
451
452
- Saha, N., & Mukul, S. A. (2022). Visitor’s willingness to pay for cultural ecosystem services in Bangladesh: An assessment for Lawachara National Park, a biodiversity hotspot. *Small-Scale Forestry*, 21, 185–201. 453
454
455
- Sarkar, P., Banerjee, S., Biswas, S., Saha, S., Pal, D., Naskar, M. K., Srivastava, S. K., Barman, D., Kar, G., & Mukul, S. A. (2024). Contribution of mangrove ecosystem services to local livelihoods in the Indian Sundarbans. *Sustainability*, 16, 6804. 456
457
458
- Schmitt, K., & Duke, N. C. (2015). Mangrove management, assessment and monitoring. In M. Köhl & L. Pancel (Eds.), *Tropical Forestry Handbook*. Springer. 459
460
- Siddiqi, N. A. (2001). *Mangrove forestry in Bangladesh*. Institute of Forestry and Environmental Science, Chittagong University. 461
462
- Spalding, M., Kainuma, M., & Collins, L. (2010). *Word Atlas of Mangroves*. Earthscan. 463
- Stubbs, B. J., & Saenger, P. (2002). The application of forestry principles to the design, execution and evaluation of mangrove restoration projects. *Bois et Forêts des Tropiques*, 273, 5–21. 464
465
- Thomas, N., Lucas, R., Bunting, P., Hardy, A., Rosenqvist, A., & Simard, M. (2017). Distribution and drivers of global mangrove forest change, 1996–2010. *PLoS One*, 12, e0179302. 466
467
- Twilley, R. R., Chen, R. H., & Hargis, T. (1992). Carbon sinks in mangroves and their implications to the carbon budget of tropical coastal ecosystems. *Water, Air, and Soil Pollution*, 64, 265–288. 468
469
- Vo, Q. T., Kuenzer, C., Vo, Q. M., Moder, F., & Oppelt, N. (2012). Review of valuation methods for mangrove ecosystem services. *Ecological Indicators*, 23, 431–446. 470
471
- Walters, B. B., Rönnbäck, P., Kovacs, J. M., Crona, B., Hussain, S. A., Badola, R., et al. (2008). Ethnobiology, socio-economics and management of mangrove forests: A review. *Aquatic Botany*, 89, 220–236. 472
473
474
- Wang, T., Zuo, Y., Manda, T., Hwarari, D., & Yang, L. (2025). Harnessing artificial intelligence, machine learning and deep learning for sustainable forestry management and conservation: Transformative potential and future perspectives. *Plants*, 14, 998. 475
476
477
- zu Ermgassen, P. S. E., Worthington, T. A., Gair, J. R., et al. (2025). Mangroves support an estimated annual abundance of over 700 billion juvenile fish and invertebrates. *Communications Earth & Environment*, 6, 299. 478
479
480