1

Invasive Alien Species of Bangladesh

Sharif A. Mukul[1,2], Mohammed Abu Sayed Arfin-Khan[3], and Mohammad Belal Uddin[3]

[1] *Tropical Forests and People Research Centre, University of the Sunshine Coast, Maroochydore, Australia*
[2] *Centre for Research on Land-use Sustainability, Dhaka, Bangladesh*
[3] *Department of Forestry and Environmental Science, School of Agriculture and Mineral Sciences, Shahjalal University of Science and Technology, Sylhet, Bangladesh*

1.1 Introduction

During the past decades, invasive alien species (IAS) have gained wider recognition by scientists and policymakers for their severe ecological and economic impacts worldwide (Early et al. 2016; Mukul et al. 2006). At the same time, IAS has been identified as one of the greatest threats to native ecosystems, habitat, and species (Guido and Pillar 2017; Mukul et al. 2006). Increasing globalization facilitates the arrival of IAS in new regions, and environmental changes, including global warming, facilitate their establishment worldwide (Early et al. 2016). Human-mediated transport beyond biogeographic barriers has also led to the introduction and establishment of IAS in new regions (Dawson et al. 2017; Turbelin et al. 2017). IAS recognized for their rapid growth, efficient dispersal capabilities, large reproductive output, and tolerance to a broad range of environmental conditions over native or local species (Campbell 2005; Hellmann et al. 2008).

Bangladesh has a long history of species introduction from different countries or geographic regions (Barua et al. 2001; Hossain 2009). More than 300 exotic species are assumed to be either domesticated or cultivated in the country at different times, of which there are no records (Ali 1991; Hossain and Pasha 2001). Several plant species, mainly herbs and shrubs, were introduced in the country during the British colonial period for their aesthetic value (Mukul et al. 2006). Most of the timber species were introduced during the late 1980s to meet the country's rapidly growing demand for timber, fuelwood, and fodder (Ahmed et al. 2008; Mukul et al. 2006). In recent years, some fish species have been introduced primarily to increase productivity to support the nutrients needs of a huge population (Galib and Mohsin 2010; Hossain 2009). Although IAS is a growing reality in Bangladesh, they are still largely ignored and unnoticed (Akter and Zuberi 2009). This chapter aims to provide an overview of IAS in Bangladesh, their impacts on ecosystems and society,

Invasive Alien Species: Observations and Issues from Around the World, Volume 2: Issues and Invasions in Asia and the Pacific Region, First Edition. Edited by T. Pullaiah and Michael R. Ielmini.
© 2021 John Wiley & Sons Ltd. Published 2021 by John Wiley & Sons Ltd.

introduction pathways, and strategies for management with a general background of the country's geography, land-use/land cover, ecosystem, and biodiversity.

1.2 Geography and Climate of Bangladesh

Bangladesh is a country of about 143 999 square kilometers (km^2) and is bordered by India in the west, north, and east except for a small portion in the southeast by Myanmar (Figure 1.1). The entire south is occupied by the Bay of Bengal. Geographically, the country is located between 88° 01′E and 92° 41′ East longitude and between 20° 34′ and 26° 33′ North latitude (Rashid 1991). The majority of the country's land is formed by river alluvium from the Ganges and the Brahmaputra and their tributaries, which consists mostly of flood plains (80%) with some hilly areas (12%) (Sohel et al. 2015). More than 50 transboundary rivers feed into the country, effectively creating the world's second largest riverine drainage basin, the Ganges–Brahmaputra–Meghna (GBM) basin, and the largest delta of the world (Hofer and Messerli 2006).

Bangladesh enjoys a tropical monsoon climate, which is characterized by high temperatures, heavy rainfall, excessive humidity, and fairly marked seasonal variations (Brammer 2016). There are three distinct seasons in the country: a hot, humid summer from March to June; a cool, rainy monsoon season from June to October; and a cool, dry winter from October to March. Maximum summer temperatures range between 30 and 40 °C. April is the warmest month, while January is the coldest month with an average temperature of about 10 °C all over the country. Much of the country receives at least 2000 mm of annual rainfall, while part of the west, northwest, and north-eastern Bangladesh receive the greatest average precipitation, often over 4000 mm per year (Rashid 1991).

1.3 Land-Use/Land Cover and Ecosystems of Bangladesh

In Bangladesh, agriculture is the major land-use, comprising nearly 65% of the country's total land area, followed by forests (17.5%), urban/built-up areas (8%), and water (Figure 1.2; Mukul et al. 2014). Forests occupy nearly 2.53 million hectares in Bangladesh and mostly degraded in nature (Mukul et al. 2017). Of these, 1.52 million hectares are under the jurisdiction of the Bangladesh Forest Department (FD) (Mukul et al. 2017; Table 1.1). The remaining forests areas are unclassed state forests under the control of district administration and homestead forests owned by smallholder landowners (Khan et al. 2007; Mukul 2007).

Majority of the lands are arable in Bangladesh, therefore, suitable for agriculture. Among the field crops, rice is the major staple crop, occupying 70% of the gross cropped area under agricultural land-use. Agricultural land often subjected to weed and insect attack influencing productivity and management costs. The country's water body mainly comprised of rivers, ponds, canals, and lakes filled with a variety of aquatic plants, fish, and migratory birds. Among the forests, mangrove forests in coastal regions of the country are most diverse and in relatively good condition (Mukul et al. 2014). People are highly dependent on hill forests and Sal forests and FD established several plantations using native and exotic species there (Khan et al. 2007).



Figure 1.1 Major land-use/land cover and ecosystem types of Bangladesh. *Source*: Bangladesh Forest Department (2016).



Figure 1.2 Major land-use/land cover of Bangladesh.

Table 1.1 Forest ecosystems of Bangladesh.

Forest type	Area (million ha)	Percentage of country's land area (%)
Hill Forests	0.67	4.5
Sal Forests	0.12	0.8
Mangrove Forests	0.60	4.1
Mangrove Plantations	0.13	0.9
Unclassed State Forest	0.73	4.9
Homestead Forests	0.27	1.8
Total	2.53	17.5

Source: Mukul SA, Khan MASA, Uddin MB (2020). Invasive alien species of Bangladesh. Preprints 2020, 2020020357.

1.4 Biodiversity in Bangladesh

Although a small country, Bangladesh is exceptionally endowed with a rich biodiversity due to its unique geophysical location (Mukul et al. 2018). An estimated 5700 species of angiosperms alone, including 68 woody legumes, 130 fiber yielding plants, 500 medicinal

Table 1.2 Faunal diversity (vertebrates) of Bangladesh.

Group	Total no. of species	Extinct	Threatened			
			Critically endangered	Endangered	Vulnerable	Total
Amphibians	49	0	2	3	5	10
Reptiles	167	1	17	10	11	39
Birds	566	19	10	12	17	58
Mammals	138	11	17	12	9	49
Total	920	31	46	37	42	156

Source: IUCN (2015).

plants, 29 orchids, 3 gymnosperms, and 1700 pteridophytes have been recorded from the country (Islam 2003). The country also possesses a rich faunal diversity although many are now under different degrees of threats due to changing climate and anthropogenic pressure (Alamgir et al. 2015; Mukul et al. 2012, 2019). Approximately 138 species of mammals, 566 species of birds, 167 species of reptiles, 49 species of amphibians, 708 species of marine and freshwater fish, 2493 species of insects, 19 species of mites, 164 species of algae (and seaweed), and 4 species of echinoderms have so far been recorded from the country (Table 1.2) (IUCN 2015).

1.5 Invasive Alien Species in Bangladesh

A complete and reliable inventory of IAS is still lacking in Bangladesh. The majority of the exotic species were introduced into the country without any proper documentation (Barua et al. 2001). There are also contradictions in the definition and the use of the term IAS in Bangladesh and not all the exotics are harmful (Mukul et al. 2006). Table 1.3 provides a list of common IAS in Bangladesh based on secondary literature. Most of the IAS in the country were brought by settlers, invaders, seamen, and traders. Few of the invasive species, mainly plants, were introduced during the colonial period as an ornamental species for their aesthetic value (Mukul et al. 2006). Perhaps the first widely introduced IAS in Bangladesh is Water Hyacinth (*Eichhornia crassipes*) which was brought from Brazil during the British colonial period (Barua et al. 2001). In recent years, deliberate preferences of fast-growing, high-yielding cultivars over native species caused the introduction of some potentially harmful exotic species in the country (Hossain 2009). The introduction of Acacia and Eucalyptus from Australia during the 1980s created several controversies in the country. All the species belonging to these two genera are proven to be rivals to the endemic flora and found to be environmentally unfriendly (Ameen 1999). The initial list of IAS prepared by Zabala (1990) of angiospermic flora of the country listed nine species, i.e. *Acacia auriculiformis*, *A. mangium*, *Albizia falcataria*, *Dalbergia sissoo, Eucalyptus camaldulensis, E. brassiana, Leucaena leucocephala, Swietenia macrophylla,* and *S. mahagony,* although not all of them fulfill the criteria of IAS (Akter and Zuberi 2009).

Other invasive plants are widespread in forests, wastelands, agricultural landscapes, and aquatic ecosystems mostly as noxious weed (see Akter and Zuberi 2009; Barua et al. 2001;

Table 1.3 Invasive alien species reported from different ecosystems of Bangladesh.

Life form	Scientific name	Common name	Native range	Habitat type(s)	Major use(s)	Occurrence	Level of Invasion
Bird	*Columba livia*	Rock pigeon	Europe	TER	P	++	++
Fish	*Aristichthys nobilis*	Bighead carp	China	AQU	F	++	++
	Carassius auratus	Goldfish	Europe, Asia	AQU	O	+	+
	Clarias gariepinus	African sharptooth catfish	Africa	AQU	F	++	++
	Ctenopharyngodon idella	Grass carp	China, Russia	AQU	F	++	++
	Cyprinus carpio	Common carp	Africa, China, Russia	AQU	F	++	++
	Gambusia affinis	Mosquito fish	Armenia, Mexico	AQU	O	++	++
	Hypophthalmichthys molitrix	Sliver carp	China, Russia	AQU	F	++	++
	Lebistes reticulatus	Guppy	South America	AQU	F	++	++
	Mylopharyngodon piceus	Black carp	China	AQU	F	++	++
	Oreochromis mossambicus	Common tilapia	Africa	AQU	F	++	++
	Oreochromis niloticus	Nilotica	Africa	AQU	F	++	++
	Pangasius giganticus	Giant pangas	NA	AQU	F	++	++
	Pangasius sutchi	Pangas	Thailand	AQU	F	++	++
	Pterygoplichthys multiradiatus	Armored catfish	NA	AQU	O	++	++
	Puntius gonionotus	Thai sarpunti	Thailand	AQU	F	++	++
	Trichogaster pectoralis	Siamese gourami	Singapore	AQU	F, O	++	++
Insect	*Bemisia tabaci*	Silverleaf whitefly	NA	AGR	P	++	++
	Diaphorina citri	Asian citrus psyllid	Asia	AGR	P	++	++
	Paratrechina longicornis	Crazy ant	Africa	FOR	P	++	++

	Tapinoma melanocephalum	Ghost ant	NA	FOR	P	++	++
	Trogoderma granarium	Khapra beetle	India	AGR	P	++	++
Mollusk	*Achatina fulica*	Giant African land snail	Africa	TER	P	++	+
Plant	*Acacia auriculiformis*	Acacia	Australia and Pacific	FOR	Ti, Ni	P	+
	Acacia mangium	Black wattle	Australia and Pacific	FOR	Ti, Ni	P	+
	Acanthospermum hispidum	Hispid starburr	Central and South America	AGR, WF	W, M	++	+
	Ageratum conyzoides	Billy goat weed	South America	FOR, AGR, WF	W, M	++	++
	Alternanthera ficoidea	Joseph's coat	South America	WF	W	++	+
	Atylosia scarabaeoides	Wild pigeonpea	Australia	AGR, WF	W	++	+
	Cardamine flexuosa	Wavy bittercress	Europe	AGR, WF	W	++	++
	Cassia occidentalis	Mogdad coffee	South America	F, WF	W, M	++	+
	Cestrum diurnum	Day-blooming cestrum	South America	AGR, WF	O	+	+
	Chromolaena odorata	Siam weed	South America	FOR, AGR	W, M	++	++
	Commelina obliqua	Day flower	Java	WF	W, Fo	++	+
	Convolvulus arvensis	Binweed	Europe	WF	W	++	+
	Croton bonplandianum	Croton	South America	AGR, WF	W, M	++	+
	Cyperus rotundus	Nut grass	Africa, Europe	AGR, WF	W	++	++
	Dalbergia sissoo	Sissoo	India	FOR		P	?
	Eichhornia crassipes	Water hyacinth	South America	AQU	W	++	++

(*Continued*)

Table 1.3 (Continued)

Life form	Scientific name	Common name	Native range	Habitat type(s)	Major use(s)	Occurrence	Level of Invasion
	Eucalyptus camaldulensis	Eucalyptus	Australia	FOR (P)	Ti	P	+
	Eucalyptus tereticornis	Eucalyptus	Australia	FOR	Ti	P	?
	Eucalyptus brassiana	Eucalyptus	Australia	FOR	Ti	P	?
	Evolvulus nummularius	Roundleaf blindweed	West Indies	AGR, WF	W	++	+
	Falcataria moluccana	Malacana koroi	Australia and Pacific	FOR	Ti, Ni	P	+
	Hyptis suaveolens	Pignut	South America	WF	W, M	++	+
	Imperata cylindrica	Cogon grass	North America	FOR, WF	W	++	++
	Ipomoea carnea	Pink morning glory	South America	AQU	W, M	++	++
	Lantana camara	Lantana	South America	FOR	W, O	++	++
	Leucaena leucocephala	Wild tamarind	South America	FOR	Fo, Ni	++	++
	Limnocharis flava	Yellow burhead	South America	AQU	W	++	++
	Ludwigia adscendens	Water primrose	Central America	AQU, AGR	W	++	+
	Mikania micrantha	Mile-a-minute weed	South America	FOR, AGR, WF	W, M	++	++
	Mimosa pudica	Touch-me-not	South America	FOR, AGR, WF	W, M	++	++
	Melastoma malabathricum	Indian rhododendron	India, Japan, Australia	FOR, WF	W, M	++	+
	Parthenium hysterophorus	Carrot grass	South America	WF	W	++	++
	Pinus caribaea	Caribaea pine	Central America	FOR	Ti, O	P	+
	Pinus oocarpa	Pine	Central America	FOR	Ti, O	P	?

Swietenia mahagoni	True mahogani	North America and Caribbean	FOR	Ti	P	?
Swietenia macrophylla	Mahogani	Central and South America	FOR (P)	Ti	P	?
Salvinia molesta	Giant salvinia	South America	AQU	W	++	++
Tectona grandis	Teak	Myanmar, India	FOR	Ti	P	+
Urena lobata	Caesar weed	South America, Africa	FOR, WF	W	++	++
Xylia dolabriformis	Pynkado	Myanmar, India	FOR	Ti	P	?

Habitat type: TER – terrestrial, AQU – aquatic, AGR – agricultural, FOR – forest, WF – waste and fallow land;
Major use(s): F – food, Fo – fodder, Fu – fuel, M – medicinal, Ni – nitrogen fixation, O – ornamental, P – pest, W – weed, T – timber;
Occurrence: + – common, ++ – very common, – - rare, P – planted;
Level of invasion: + – invasive, ++ – highly invasive, ? – potentially invasive.
Source: Information for preparing this table has been taken from Akter and Zuberi (2009); Barua et al. (2001); Biswas et al. (2007); GISD (2019); Hossain (2009); Hossain and Pasha (2001); Khan et al. (2011); Mukul et al. (2006); Uddin et al. (2013).

Biswas et al. 2007, 2012; Hossain 2009; Hossain and Pasha 2001; Islam et al. 2003; Khan et al. 2011; Mukul et al. 2006; Uddin et al. 2013). Some of the world's worst IAS like, *Chromolaena odorata, Lantana camara,* and *Mikania micrantha,* are also abundant in Bangladesh with a detrimental impact on local species and natural ecosystems (Ahmed et al. 2007). Figure 1.3 shows some of the common invasive alien plant species in different ecosystems of the country.

Among the fauna, most of the reported IAS are fish and mostly carps. The predatory habit of these carp species is well recognized. Although others are not predatory, their fecundity and growth rate are extremely high, and they are able to breed naturally (Barua et al. 2001). Many of these species were brought to the country from Thailand during the 1990s. Currently, the most "disastrous" invasive alien fish species in the country are – *Clarias gariepinus*, *Pangasius sutchi*, *Pangasius giganticus*, *Tilapia mossambica,* and *Oreochromis niloticus* (Rahman 1997). All of these species also recognized as the world's worst IAS (Lowe et al. 2000). A large number of ornamental and game fish have also been introduced in Bangladesh in the past decades (Galib and Mohsin 2010).

Other invasive alien fauna includes a bird species (*Columba livia*), several insects, and a species of mollusks (*Achatina fulica*). This list, however, does not epitomize the actual situation of IAS in the country. Many agricultural pests and microorganisms which have been brought to the country during agricultural trade remain unrecorded. In recent years,



Figure 1.3 Some common invasive alien plant species in Bangladesh (clockwise) – (a) Touch-me-not (*Mimosa pudica*), (b) Water hyacinth (*Eichhornia crassipes*), (c) Billy goat weed (*Ageratum conyzoides*), and (d) Lantana (*Lantana camara*). Photo: Sharif A. Mukul.

unregulated wildlife trade and imports, and a growing urban market of wildlife also caused the introduction of some potentially invasive fauna into the country's terrestrial ecosystem many of which remained unidentified.

1.6 Threats from Invasive Alien Species in Bangladesh

IAS in natural ecosystems is one of the major sources of biodiversity loss worldwide (Guido and Pillar 2017). They also disrupt native ecosystems, degrade wild habitats, and jeopardize endangered plants and animals (Early et al. 2016; Padmanaba and Corlett 2014). In natural ecosystems, the introduction of some IAS at the initial stage may although enhance primary production, some ecosystem processes (e.g. N-cycling) and services, in successive rotations, there might be a sharp fall in those services and benefits (Al-Mamun et al. 2013; Vila et al. 2011). Many invasive species also possess broad climatic tolerances and large geographic ranges, which positively influence their ability to adapt and reproduce over native species (Hellmann et al. 2008; Wittenberg and Cock 2001).

The economic loss due to IAS in the agricultural sector is well documented not only in Bangladesh but also globally. The cost associated with eradication and management of IAS in agricultural systems is huge. In forest ecosystems, fast-growing exotic species like *Acacia* sp., *Eucalyptus* sp., and *Leucaena leucocephala* compete with native species for nutrition and water from the soil. Many of them also hold an allelopathic effect and suppress the development of undergrowth. Other IAS, like the Common Water Hyacinth (*Eichhornia crassipes*) is responsible for the large-scale obstruction of the navigation system in inland water bodies (Figure 1.4), negatively affecting fish production and disrupt aquatic

Figure 1.4 Common Water Hyacinth (*Eichhornia crassipes*) obstructing local water bodies in Bangladesh. Photo: Sharif A. Mukul.

ecosystem. The feeding habit of introduced carp species in the country frequently stated for pond bank erosion, increased turbidity, and elevated nutrient concentrations. These alterations to physical and chemical conditions also have ecological consequences, such as increased phytoplankton density in response to elevated nutrient levels. These carp species compete with native fish species for natural foods, which ultimately make the local species vulnerable. Prolific breeding of species, like Tilapia (*Oreochromis mossambicus*) makes natural food unavailable to local fish species. The African Sharptooth Catfish (*Clarias gariepinus*) directly feed on native fish species thereby causing their depletion in inland water bodies.

The impacts of IAS on a country's genetic pool are also significant. IAS are known to hybridize with native species, thus altering genetic diversity and integrity of native species. Exotic species are also common pathways of transmitting disease and pathogens in the country. For example, the invasive *Argulus* sp. has been introduced with the introduction of carps in the country. It is also widely believed that the white spot syndrome virus (WSSV) of Shrimp which caused a devastating fall in shrimp production and export from Bangladesh was introduced from the different stock of fish imported from neighboring countries.

1.7 Conclusion and Recommendations

A comprehensive assessment and framework for identifying IAS are still lacking in Bangladesh. The country's extended borders and ports are largely unprotected. Enforcement of existing safeguard mechanisms such as proper quarantine measures while importing any exotic species into the country is also weak. Conservation of biodiversity, however, is a national priority of the country, as a signatory to the United Nations Conventions on Biological Diversity (UNCBD). A statutory body consisting of local experts and standard and comprehensible procedures for the introduction, monitoring, and management of alien species in country's agriculture, forestry, and fisheries sector is indispensable. The government should also be cautious while introducing new species in the country with clear and effective quarantine regulations.

While mechanical, chemical, and biological control is the widely used approach for controlling IAS in any invaded ecosystem, they require skilled manpower, technology, and expertise and could be extremely costly and labor-intensive. Preventive measures, therefore, should be the highest priority of the country. Appropriate policies and rules must be developed to facilitate the control of IAS in the country. Community awareness and local people's involvement with relevant stakeholders should be advocated for managing IAS in country's different land-use/land cover and ecosystems. Community-based vigilance and monitoring for known pathways of introduction of IAS should also be undertaken as much as possible. In the case of transboundary landscapes and water bodies, regional cooperation must be sought for the effective management of IAS. Last but not the least, capacity building of government staffs involved in quarantine and border control must be guaranteed on a regular basis.

References

Ahmed, R., Uddin, M.B., Khan, M.A.S.A. et al. (2007). Allelopathic effects of *Lantana camara* on germination and growth behavior of some agricultural crops in Bangladesh. *Journal of Forestry Research* 18: 301–304.

Ahmed, Z.U., Begum, Z.N.T., Hassan, M.A. et al. (eds.) (2008). *Encyclopedia of Flora and Fauna of Bangladesh*. Dhaka: Asiatic Society of Bangladesh.

Akter, A. and Zuberi, M.I. (2009). Invasive alien species in northern Bangladesh: identification, inventory and impacts. *International Journal of Biodiversity and Conservation* 1: 129–134.

Alamgir, M., Mukul, S.A., and Turton, S. (2015). Modelling spatial distribution of critically endangered Asian elephant and Hoolock gibbon in Bangladesh forest ecosystems under a changing climate. *Applied Geography* 60: 10–19.

Ali, S.B. (1991). Plant introduction – a continuous process. In: *Two Centuries of Plant Studies in Bangladesh and Adjacent Regions* (ed. A.K.M.N. Islam), 97–153. Dhaka: Asiatic Society of Bangladesh.

Al-Mamun, M., Poostforush, M., Mukul, S.A. et al. (2013). Biosorption of as(III) from aqueous solution by *Acacia auriculiformis* leaves. *Scientia Iranica* 20: 1871–1880.

Ameen, M. (1999). Development of Guiding Principles for the Prevention of Impacts of Alien Species. Paper presented at 'Consultative Workshop in Advance of the 4th Meeting of SBSTTA to the CBD', organised by IUCN Bangladesh at Dhaka on 25 May 1999.

Bangladesh Forest Department (2016). Map of Forests of Bangladesh. www.bforest.gov.bd (accessed 15 June 2018).

Barua, S.P., Khan, M.M.H., and Reza, A.H.M.A. (2001). The status of alien invasive species in Bangladesh and their impact on the ecosystems. In: *Alien Invasive Species – Report of the Workshop on Alien Invasive Species* (ed. P. Balakrishna), 1–7. Colombo, Sri Lanka: IUCN Regional Biodiversity Programme of Asia.

Biswas, S.R., Choudhury, J.K., Nishat, A., and Rahman, M.M. (2007). Do invasive plants threaten the Sundarbans mangrove forest of Bangladesh? *Forest Ecology and Management* 245: 1–9.

Biswas, S.R., Khan, M.S.I., and Mallik, A.U. (2012). Invaders' control on post-disturbance succession in coastal mangroves. *Journal of Plant Ecology* 5: 157–166.

Brammer, H. (2016). *Bangladesh: Landscapes, Soil Fertility and Climate Change*. Dhaka: University Press Limited.

Campbell S (2005) A global perspective on forest invasive species: the problem, causes, and consequences. In: McKenzie P, Brown C, Jianghua S, Jian W (eds.), *The unwelcome guests – Proceedings of the Asia-Pacific Forest Invasive Species Conference*, FAO-Regional office for Asia and the Pacific, Bangkok, Thailand, pp. 9–10.

Dawson, W., Moser, D., van Kleunen, M. et al. (2017). Global hotspots and correlates of alien species richness across taxonomic groups. *Nature Ecology & Evolution* 1: 0186.

Early, R., Bradley, B.A., Dukes, J.S. et al. (2016). Global threats from invasive alien species in the twenty-first century and national response capacities. *Nature Communications* 7: 12485.

Galib, S.M. and Mohsin, A.B.M. (2010). Exotic ornamental fishes of Bangladesh. *Bangladesh Journal of Progressive Science and Technology* 8: 255–258.

Global Invasive Species Database (GISD) (2019). Bangladesh country profile. http://issg.org/database/species/search.asp?sts=sss&st=sss&fr=1&x=0&y=0&sn=&rn=Bangladesh&hci=-1&ei=-1&lang=EN (accessed 1 December 2019).

Guido, A. and Pillar, V.D. (2017). Invasive plant removal: assessing community impact and recovery from invasion. *Journal of Applied Ecology* 54: 1230–1237.

Hellmann, J.J., Byers, J.E., Bierwagen, B.G., and Dukes, J.S. (2008). Five potential consequences of climate change for invasive species. *Conservation Biology* 22: 534–543.

Hofer, T. and Messerli, B. (2006). *Floods in Bangladesh: History, Dynamics and Rethinking the Role of the Himalayas*. Tokyo, Japan: United Nations University Press.

Hossain, M.K. (2009). Alien invasive plant species and their effects on hill forest ecosystems of Bangladesh. In: *Invasive Plants and Forest Ecosystems* (eds. R.K. Kohli, S. Jose, H.P. Singh and D.R. Batish), 133–141. New York;: CRC Press.

Hossain, M.K. and Pasha, M.K. (2001). Alien invasive plants in Bangladesh and their impacts on the ecosystem. In: *Assessment and Management of Alien Species that Threatened Ecosystem, Habitats and Species*, CBD Technical Paper No. 21, 73–75. Montreal, Canada: Secretariat of the Convention on Biological Diversity.

Islam, S.S. (2003). *State of forest genetic resources conservation and management in Bangladesh*. Forest Genetic Resources Working Papers, Working Paper FGR/68E. Forest Resources Development Service, Forest Resources Division, FAO, Rome, Italy.

Islam, M.M., Amin, A.S.M.R., and Sarker, S.K. (2003). National report on alien invasive species of Bangladesh. In: *Invasive Alien Species in South-Southeast Asia: National Reports & Directory of Resources* (eds. N. Pallewatta, J.K. Reaser and A.T. Gutierrez), 7–24. Cape Town, South Africa: Global Invasive Species Programme.

IUCN (2015). *Red List of Bangladesh – A Brief on Assessment Result 2015*. Dhaka, Bangladesh: The World Conservation Union.

Khan, M.A.S.A., Uddin, M.B., Uddin, M.S. et al. (2007). Distribution and status of forests in the tropic: Bangladesh perspective. *Proceedings of the Pakistan Academy of Sciences* 44: 145–153.

Khan, M.A.S.A., Sultana, F., Rahman, M.H. et al. (2011). Status and ethno-medicinal usage of invasive plants in traditional health care practices: a case study from northeastern Bangladesh. *Journal of Forestry Research* 22: 649–658.

Lowe, S., Brown, M., Boudjelas, S., and Poorter, M.D. (2000). *100 of the World's Worst Invasive Alien Species: A Selection from the Global Invasive Species Database*. Auckland, New Zealand: The Invasive Species Specialist Group (ISSG) of Species Survival Commission of IUCN.

Mukul, S.A. (2007). Biodiversity conservation strategies in Bangladesh: the state of protected areas. *Tigerpaper* 34: 28–32.

Mukul, S.A., Uddin, M.B., and Tito, M.R. (2006). Study on the status and various uses of invasive alien plant species in and around Satchari National Park, Sylhet, Bangladesh. *Tigerpaper* 33: 28–32.

Mukul, S.A., Rashid, A.Z.M.M., and Uddin, M.B. (2012). The role of spiritual beliefs in conserving wildlife species in religious shrines of Bangladesh. *Biodiversity* 13: 108–114.

Mukul, S.A., Biswas, S.R., Rashid, A.Z.M.M. et al. (2014). A new estimate of carbon for Bangladesh forest ecosystems with their spatial distribution and REDD+ implications. *International Journal of Research on Land-use Sustainability* 1: 33–41.

Mukul, S.A., Rashid, A.Z.M.M., and Khan, N.A. (2017). Forest protected area systems and biodiversity conservation in Bangladesh. In: *Protected Areas: Policies, Management and*

Future Directions (eds. S.A. Mukul and A.Z.M.M. Rashid), 157–177. USA: Nova Science Publishers.

Mukul, S.A., Biswas, S.R., and Rashid, A.Z.M.M. (2018). Biodiversity in Bangladesh. In: *Global Biodiversity*, Selected Countries in Asia, vol. 1 (ed. T. Pullaiah), 93–107. Canada: Apple Academic Press.

Mukul, S.A., Alamgir, M., Sohel, M.S.I. et al. (2019). Combined effects of climate change and sea-level rise project dramatic habitat loss of the globally endangered Bengal tiger in the Bangladesh Sundarbans. *Science of the Total Environment* 663: 830–840.

Mukul SA, Mohammed Abu Sayed Arfin K, Uddin MB (2020). Invasive alien species of Bangladesh. Preprints 2020, 2020020357.

Padmanaba, M. and Corlett, R.T. (2014). Minimizing risks of invasive alien plant species in tropical production forest management. *Forests* 5: 1982–1998.

Rahman, A.K.A. (1997). Fish marketing in Bangladesh. In: *Openwater Fisheries of Bangladesh* (eds. C.F. Tsai and M.Y. Ali). Dhaka: University Press Limited.

Rashid, H.E. (1991). *Geography of Bangladesh*. Dhaka: University Press.

Sohel, M.S.I., Mukul, S.A., and Chicharo, L. (2015). A new ecohydrological approach for ecosystem service provisions and sustainable management of aquatic ecosystems in Bangladesh. *Ecohydrology & Hydrobiology* 15: 1–12.

Turbelin, A.J., Malamud, B.D., and Francis, R.A. (2017). Mapping the global state of invasive alien species: patterns of invasion and policy responses. *Global Ecology and Biogeography* 26: 78–92.

Uddin, M.B., Steinbauer, M.J., Jentsch, A. et al. (2013). Do environmental attributes, disturbances, and protection regimes determine the distribution of exotic plant species in Bangladesh forest ecosystem? *Forest Ecology and Management* 303: 72–80.

Vila, M., Espinar, J.L., Hejda, M. et al. (2011). Ecological impacts of invasive alien plants: a meta-analysis of their effects on species, communities and ecosystems. *Ecology Letters* 14: 702–708.

Wittenberg, R. and Cock, M.J.W. (2001). *Invasive Alien Species: A Toolkit of Best Prevention and Management Practices*. Wallingford, UK: CAB International.

Zabala, N.Q. (1990). *Silviculture of Species*. Bangladesh: Institute of Forestry and Environmental Science, Chittagong University.